



TITLE:

NOTES ON CERTAIN CLASSES OF ANALYTIC FUNCTIONS DEFINED BY RUSCHEWEYH DERIVATIVES(Topics in Univalent Functions and Its Applications)

AUTHOR(S):

CHEN, MING-PO; YU, CHING-SHU; OWA, SHIGEYOSHI

CITATION:

CHEN, MING-PO ...[et al]. NOTES ON CERTAIN CLASSES OF ANALYTIC FUNCTIONS DEFINED BY RUSCHEWEYH DERIVATIVES(Topics in Univalent Functions and Its Applications). 数理解析研究所講究録 1990, 714: 73-85

ISSUE DATE:

1990-03

URL:

<http://hdl.handle.net/2433/101734>

RIGHT:

NOTES ON CERTAIN CLASSES OF ANALYTIC FUNCTIONS DEFINED BY RUSCHEWEYH DERIVATIVES

MING-PO CHEN (台湾中央研究院)

CHING-SHU YU (中原大学)

SHIGEYOSHI OWA (近畿大学理工・尾和重義)

1. INTRODUCTION

Let A_p denote the class of functions of the form

$$(1.1) \quad f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n \quad (p \in \mathbb{N} = \{1, 2, 3, \dots\})$$

which are analytic in the unit disk $U = \{z: |z| < 1\}$. For functions

$$(1.2) \quad f_j(z) = z^p + \sum_{n=p+1}^{\infty} a_{n,j} z^n \quad (j = 1, 2)$$

belonging to Λ , we denote by $f_1 * f_2(z)$ the convolution of $f_1(z)$ and $f_2(z)$, that is,

$$(1.3) \quad f_1 * f_2(z) = z^p + \sum_{n=p+1}^{\infty} a_{n,1} a_{n,2} z^n.$$

1980 Mathematics Subject Classification (1985 Revision). Primary 30C45.

Using the convolution, we define the Ruscheweyh derivative $D^{\alpha+p-1}f(z)$ by

$$(1.4) \quad D^{\alpha+p-1}f(z) = \frac{z^p}{(1-z)^{\alpha+p}} * f(z) \quad (\alpha > -p)$$

for $f(z) \in A_p$. A function $f(z)$ belonging to A_p is said to be in the class $R(\alpha+p-1)$ if it satisfies

$$(1.5) \quad \operatorname{Re} \left\{ \frac{D^{\alpha+p}f(z)}{D^{\alpha+p-1}f(z)} \right\} > \frac{\alpha+p-1}{\alpha+p} \quad (z \in U),$$

where $\alpha > -p$.

For an integer α greater than $-p$, Soni [4] proved

THEOREM A. $R(\alpha+p) \subset R(\alpha+p-1)$.

for $\alpha = n \in N_0 = \{0, 1, 2, \dots\}$ and $p = 1$, Singh and Singh [3] proved

THEOREM B. $R(n+1) \subset R(n)$.

In the present paper, we improve the above results.

2. PROPERTIES OF THE CLASS $R(\alpha+p-1)$

We begin with the statement of the following lemma due to Miller and

Mocanu [2].

LEMMA 1. Let $\varphi(u, v)$ be a complex valued function,

$$\varphi: D \rightarrow \mathbb{C}, D \subset \mathbb{C}^2 \quad (\mathbb{C} \text{ is the complex plane}),$$

and let $u = u_1 + iu_2, v = v_1 + iv_2$. Suppose that the function $\varphi(u, v)$ satisfies

- (i) $\varphi(u, v)$ is continuous in D ;
- (ii) $(1, 0) \in D$ and $\operatorname{Re}\{\varphi(1, 0)\} > 0$;
- (iii) for all $(iu_2, v_1) \in D$ such that $v_1 \leq -\frac{1+u_2^2}{2}$,
 $\operatorname{Re}\{\varphi(iu_2, v_1)\} \leq 0$.

Let $q(z) = 1 + q_1 z + q_2 z^2 + \dots$ be regular in U such that $(q(z), zq'(z)) \in D$ for all $z \in U$. If

$$\operatorname{Re}\{\varphi(q(z), zq'(z))\} > 0 \quad (z \in U),$$

then $\operatorname{Re}(q(z)) > 0 \quad (z \in U)$.

Applying the above lemma, we derive

THEOREM 1. If $f(z) \in R(\alpha+p)$, $\alpha \geq 1-p$, $p \in \mathbb{N}$, then

$$(2.1) \quad \operatorname{Re} \left\{ \frac{D^{\alpha+p} f(z)}{D^{\alpha+p-1} f(z)} \right\} > \beta(\alpha, p) \quad (z \in U),$$

where

$$(2.2) \quad \beta(\alpha, p) = \frac{2(\alpha+p)-3+\sqrt{4(\alpha+p)^2-4(\alpha+p)+9}}{4(\alpha+p)}.$$

PROOF. We define the function $q(z)$ by

$$(2.3) \quad \frac{D^{\alpha+p} f(z)}{D^{\alpha+p-1} f(z)} = \beta + (1-\beta)q(z),$$

where $\beta = \beta(\alpha, p)$. Noting that

$$(2.4) \quad z(D^{\alpha+p-1} f(z))' = (\alpha+p)D^{\alpha+p} f(z) - \alpha(D^{\alpha+p-1} f(z)) \quad (\alpha > -p),$$

we have, from (2.3), that

$$(2.5) \quad \frac{D^{\alpha+p+1} f(z)}{D^{\alpha+p} f(z)} = \frac{1}{\alpha+p-1} \left\{ 1 + (\alpha+p)(\beta+(1-\beta)q(z)) + \frac{(1-\beta)zq'(z)}{\beta+(1-\beta)q(z)} \right\},$$

or

$$(2.6) \quad \operatorname{Re} \left\{ \frac{D^{\alpha+p+1} f(z)}{D^{\alpha+p} f(z)} - \frac{\alpha+p}{\alpha+p+1} \right\}$$

$$= \operatorname{Re} \frac{1}{\alpha+p+1} \left\{ (1-\alpha-p) + (\alpha+p)(\beta+(1-\beta)q(z)) + \frac{(1-\beta)zq'(z)}{\beta+(1-\beta)q(z)} \right\}$$

> 0 .

Letting

$$(2.7) \quad \varphi(u, v) = \frac{1}{\alpha+p+1} \left\{ (1-\alpha-p) + (\alpha+p)(\beta+(1-\beta)u) + \frac{(1-\beta)v}{\beta+(1-\beta)u} \right\},$$

we see that

- (i) $\varphi(u, v)$ is continuous in $D = (C - \{\frac{\beta}{\beta-1}\}) \times C$;
- (ii) $(1, 0) \in D$ and $\operatorname{Re}\{\varphi(1, 0)\} = \frac{1}{\alpha+p+1} > 0$;
- (iii) for all $(iu_2, v_1) \in D$ such that $v_1 \leq -\frac{1+u_2^2}{2}$,

$$\operatorname{Re}\{\varphi(iu_2, v_1)\} = \frac{1}{\alpha+p+1} \{ (1-\alpha-p) + \beta(\alpha+p) + \frac{\beta(1-\beta)v_1}{\beta^2+(1-\beta)^2u_2^2} \}$$

$$\leq \frac{1}{\alpha+p+1} \{ (1-\alpha-p) + \beta(\alpha+p) - \frac{\beta(1-\beta)(1+u_2^2)}{2(\beta^2+(1-\beta)^2u_2^2)} \}.$$

$$= \frac{1}{\alpha+p+1} \{ 2(\alpha+p)\beta^2 - [2(\alpha+p)-3]\beta-1)(\beta+(\beta-1)u_2^2) \\$$

$$+ (1-\beta)[2(\alpha+p)(\beta-1)+1]u_2^2 \}$$

$$= \frac{(1-\beta)[2(\alpha+p)(\beta-1)+1]u_2^2}{\alpha+p+1}$$

$$\leq 0$$

because $2(\alpha+p)\beta^2 - [2(\alpha+p)-3]\beta-1 = 0$, and under the condition $\alpha+p \geq 1$, we have

$$0 < \beta \leq 1 - \frac{1}{2(\alpha+p)} < 1.$$

Therefore, the function $\varphi(u, v)$ satisfies the conditions in Lemma 1. Since $\operatorname{Re}\{q(z)\} > 0$ ($z \in U$), we conclude that

$$(2.8) \quad \operatorname{Re}\left\{\frac{D^{\alpha+p}f(z)}{D^{\alpha+p-1}f(z)}\right\} > \beta = \frac{2(\alpha+p)-3+\sqrt{4(\alpha+p)^2-4(\alpha+p)+9}}{4(\alpha+p)}$$

REMARK 1. Since

$$(2.9) \quad \operatorname{Re}\left\{\frac{D^{\alpha+p}f(z)}{D^{\alpha+p-1}f(z)}\right\} > \frac{2(\alpha+p)-3+\sqrt{4(\alpha+p)^2-4(\alpha+p)+9}}{4(\alpha+p)} > \frac{\alpha+p-1}{\alpha+p}$$

for $\alpha \geq 1-p$, Theorem 1 is the improvement of Theorem A due to Soni [4].

REMARK 2. For $\alpha = n \in N_0$ and $p = 1$, we have

$$(2.10) \quad \operatorname{Re}\left\{\frac{D^{n+1}f(z)}{D^n f(z)}\right\} > \frac{2n-1+\sqrt{4n^2+4n+9}}{4(n+1)} > \frac{n}{n+1}.$$

Therefore, Theorem 1 is also the improvement of Theorem B by Singh and Singh [3].

Taking $\alpha = 2-p$ in Theorem 1, we have

COROLLARY 1. If $f(z) \in R(2)$, then

$$(2.11) \quad \operatorname{Re}\left\{1 + \frac{zf''(z)}{f'(z)}\right\} > \frac{\sqrt{17} - 3}{4} \quad (z \in U),$$

that is, $f(z)$ is p -valently convex of order $\frac{\sqrt{17} - 3}{4}$.

PROOF. For $\alpha = 2-p$, we have

$$(2.12) \quad \operatorname{Re}\left\{\frac{D^2f(z)}{Df(z)}\right\} = \operatorname{Re}\left\{1 + \frac{1}{2} \frac{zf''(z)}{f'(z)}\right\} > \frac{1 + \sqrt{17}}{8}.$$

Therefore, (2.12) leads to

$$(2.13) \quad \operatorname{Re}\left\{1 + \frac{zf''(z)}{f'(z)}\right\} > \frac{1 + \sqrt{17}}{4} - 1 = \frac{\sqrt{17} - 3}{4}.$$

3. AN APPLICATION OF JACK'S LEMMA

In order to prove our next result, we need the following lemma by Jack [1].

LEMMA 2. Let $w(z)$ be regular in U with $w(0) = 0$. If $|w(z)|$ attains its maximum value on the circle $|z| = r$ at a point $z_0 \in U$, then

$$z_0 w'(z_0) = kw(z_0),$$

where k is real and $k \geq 1$.

THEOREM 2. If $f(z) \in A_p$ satisfies

$$(3.1) \quad \operatorname{Re} \left\{ \frac{D^{\alpha+p+1} f(z)}{D^{\alpha+p} f(z)} \right\} > \frac{1 + (\alpha+p-1)^2}{\alpha+p+1} \quad (z \in U),$$

and $\alpha + p \geq 1$ then

$$(3.2) \quad \operatorname{Re} \left[\frac{D^{\alpha+p} f(z)}{D^{\alpha+p-1} f(z)} \right]^{\frac{1}{2}} > \frac{\alpha+p-1}{\alpha+p} \quad (z \in U).$$

PROOF. For $f(z) \in A_p$ satisfying (3.1), we define the function $w(z)$ by

$$(3.3) \quad \operatorname{Re} \left[\frac{D^{\alpha+p} f(z)}{D^{\alpha+p-1} f(z)} \right]^{\frac{1}{2}} = \frac{\alpha+p-1}{\alpha+p} + \frac{1}{\alpha+p} \frac{1-w(z)}{1+w(z)} = \frac{(\alpha+p) + (\alpha+p-2)w(z)}{(\alpha+p)(1+w(z))}.$$

Then $w(z)$ is regular in U with $w(0) = 0$ and $w(z) \neq -1$ ($z \in U$).

After taking the logarithmic derivative of both sides of (3.3), using (2.4)

and (3.3) we obtain

$$(3.4) \quad \frac{D^{\alpha+p+1}f(z)}{D^{\alpha+p}f(z)} = \frac{1}{\alpha+p+1} \left\{ 1 + \left(\frac{(\alpha+p) + (\alpha+p-2)w(z)}{1+w(z)} \right)^2 - \frac{4zw'(z)}{(1+w(z))((\alpha+p) + (\alpha+p-2)w(z))} \right\}.$$

If there exists a point $z_0 \in U$ such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = 1,$$

then by Lemma 2, we have

$$z_0 w'(z_0) = kw(z_0) \quad (k \geq 1).$$

It follows from

$$\operatorname{Re} \left\{ \frac{1}{1+w(z_0)} \right\} = \frac{1}{2}$$

and

$$\operatorname{Re} \left\{ \frac{w(z_0)}{1+w(z_0)} \right\} = \frac{1}{2}$$

that

$$\begin{aligned}
 (3.5) \quad & \left\{ \frac{D^{\alpha+p+1} f(z_0)}{D^{\alpha+p} f(z_0)} \right\} \\
 = & \operatorname{Re} \frac{1}{\alpha+p+1} \left\{ 1 + \left(\frac{(\alpha+p) + (\alpha+p-2)w(z_0)}{1 + w(z_0)} \right)^2 - \frac{4kw'(z_0)}{(1+w(z_0))((\alpha+p) + (\alpha+p-2)w(z_0))} \right\}.
 \end{aligned}$$

Let $w(z_0) = e^{i\theta}$, we have

$$\begin{aligned}
 (1 + w(z_0))^2 &= (1 + e^{i\theta})^2 = (e^{-\frac{i\theta}{2}} + e^{\frac{i\theta}{2}})^2 e^{i\theta} \\
 &= (2 \cos \frac{\theta}{2})^2 e^{i\theta} = 2(1 + \cos \theta)w(z_0).
 \end{aligned}$$

It follows that

$$\begin{aligned}
 (3.6) \quad & \operatorname{Re} \left[\frac{(\alpha+p) + (\alpha+p-2)w(z_0)}{1 + w(z_0)} \right]^2 \\
 = & \operatorname{Re} \left\{ \frac{(\alpha+p)^2}{2(1+\cos \theta)e^{i\theta}} \right\} + \operatorname{Re} \left\{ \frac{2(\alpha+p)(\alpha+p-2)}{2(1+\cos \theta)} \right\} + \operatorname{Re} \left\{ \frac{(\alpha+p-2)^2 e^{i\theta}}{2(1+\cos \theta)} \right\} \\
 = & \frac{\{(\alpha+p)^2 + (\alpha+p-2)^2\} \cos \theta + 2(\alpha+p)(\alpha+p-2)}{2(1 + \cos \theta)} \\
 \leq & \frac{(\alpha+p)^2 + (\alpha+p-2)^2 + 2(\alpha+p)(\alpha+p-2)}{4} = (\alpha+p-1)^2,
 \end{aligned}$$

since $(\alpha+p)^2 + (\alpha+p-2)^2 \geq 2(\alpha+p)(\alpha+p-2)$ and $\frac{\Lambda x + B}{1+x}$ is an increasing function of x if $\Lambda \geq B$.

Next we have

$$\begin{aligned}
 (3.7) \quad & \operatorname{Re} \left\{ \frac{w(z_0)}{(1+w(z_0))((\alpha+p)+(\alpha+p-2)w(z_0))} \right\} \\
 &= \frac{1}{|(\alpha+p)+(\alpha+p-2)w(z_0)|^2} \operatorname{Re} \left\{ \frac{w(z_0)[(\alpha+p)+(\alpha+p-2)\overline{w(z_0)}]}{1+w(z_0)} \right\} \\
 &= \frac{1}{|(\alpha+p)+(\alpha+p-2)w(z_0)|^2} \left\{ (\alpha+p)\frac{1}{2} + (\alpha+p-2)\frac{1}{2} \right\} \geq 0.
 \end{aligned}$$

From (3.5), (3.6) and (3.7) we obtain

$$\operatorname{Re} \left\{ \frac{D^{\alpha+p+1}f(z_0)}{D^{\alpha+p}f(z_0)} \right\} \leq \frac{1+(\alpha+p-2)^2}{\alpha+p+1}.$$

This contradicts our condition (3.1). Therefore, we conclude that $w(z)$ satisfies $|w(z)| < 1$ for all $z \in U$, which completes the proof of our theorem.

Letting $\alpha = 2-p$, Theorem 2 gives

COROLLARY. If $f(z) \in A_p$ satisfies

$$(3.8) \quad \operatorname{Re} \left\{ \frac{D^3 f(z)}{D^2 f(z)} \right\} > \frac{2}{3} \quad (z \in U),$$

then

$$(3.9) \quad \operatorname{Re} \sqrt{2 + \frac{z f'''(z)}{f'(z)}} > \frac{\sqrt{2}}{2} \quad (z \in U).$$

ACKNOWLEDGMENT. This research of the authors was completed at Institute of Mathematics, Academia Sinica, Taipei, Taiwan while the third author was visiting from Kinki University, Higashi-Osaka, Osaka 577, Japan.

REFERENCES

- [1] I. S. Jack, Functions starlike and convex of order α , J. London Math. Soc. (2)3(1971), 469–474.
- [2] S. S. Miller and P. T. Mocanu, Second order differential inequalities in the complex plane, J. Math. Anal. Appl. 65(1978), 289–305.
- [3] R. Singh and S. Singh, Integrals of certain univalent functions, Proc. Amer. Math. Soc. 77(1979), 336–340.
- [4] A. K. Soni, Generalizations of p -valent functions via the Hadamard product, Internat. J. Math. Sci. 5(1982), 289–299.

Institute of Mathematics

Academia Sinica

Nankang, Taipei, Taiwan 11529

Republic of China

Department of Mathematics

Chung Yuan University

Chung-Li, Taiwan

Republic of China

Department of Mathematics

Kinki University

Higashi-Osaka, Osaka 577

Japan